

Computation of Koszul homology and application to partial differential systems

C. Chenavier, T. Cluzeau, A. Quadrat

8th IFAC Symposium on Systems Structure and Control

September 27, 2022

I. Motivations

- ▷ formal theory of PDEs systems
- ▷ formal integrability
- ▷ 2-acyclicity and Koszul homology

II. Computation of Koszul homology

- ▷ symbol module
- ▷ computation of homology groups
- ▷ 2-acyclicity criterion

III. Examples

- ▷ constant coefficients
- ▷ parametric coefficients
- ▷ polynomial coefficients

IV. Conclusion and perspectives

I. MOTIVATIONS

Consider a PDEs system

$$(\Sigma) : F^p \left(x^i, \frac{\partial^{|\mu|} u^j}{\partial x^\mu} \right) = 0$$

→ $x^1, \dots, x^n, u^1, \dots, u^p$: (in)dependent variables

→ F^1, \dots, F^q : equations of the system

Consider a PDEs system

$$(\Sigma) : F^p \left(x^i, \frac{\partial^{|\mu|} u^j}{\partial x^\mu} \right) = 0$$

→ $x^1, \dots, x^n, u^1, \dots, u^p$: (in)dependent variables

→ F^1, \dots, F^q : equations of the system

Formal theory

Geometric methods

- integral manifolds
- differential forms
- jet bundles

Algebraic methods

- differential elimination
- integrability conditions
- homological algebra

Consider a PDEs system

$$(\Sigma) : F^p \left(x^i, \frac{\partial^{|\mu|} u^j}{\partial x^\mu} \right) = 0$$

→ $x^1, \dots, x^n, u^1, \dots, u^p$: (in)dependent variables

→ F^1, \dots, F^q : equations of the system

Formal theory

Geometric methods

- integral manifolds
- differential forms
- jet bundles

Algebraic methods

- differential elimination
- integrability conditions
- homological algebra

Consider a PDEs system

$$(\Sigma) : F^p \left(x^i, \frac{\partial^{|\mu|} u^j}{\partial x^\mu} \right) = 0$$

→ $x^1, \dots, x^n, u^1, \dots, u^p$: (in)dependent variables

→ F^1, \dots, F^q : equations of the system

Formal theory

Geometric methods

- integral manifolds
- differential forms
- jet bundles

Algebraic methods

- differential elimination
- integrability conditions
- homological algebra

Nonlinear control problems

A fundamental notion: formal integrability

→ derivatives of F^p 's do not bring new integrability conditions

A fundamental notion: **formal integrability**

→ derivatives of F^p 's do not bring new integrability conditions

A fundamental notion: **formal integrability**

→ derivatives of F^p 's do not bring new integrability conditions

Examples

$$(\Sigma_1): \dot{u} = u$$

→ (Σ_1) is formally integrable with formal power series solutions

$$u_0 + u_0 t + \frac{u_0}{2} t^2 + \frac{u_0}{3!} t^3 + \dots$$

A fundamental notion: **formal integrability**

→ derivatives of F^p 's do not bring new integrability conditions

Examples

$$(\Sigma_1): \dot{u} = u$$

→ (Σ_1) is formally integrable with formal power series solutions

$$u_0 + u_0 t + \frac{u_0}{2} t^2 + \frac{u_0}{3!} t^3 + \dots$$

$$(\Sigma_2): u_{33} = u_{13}, \quad u_{23} = u_{13}, \quad u_{12} = u_{11}$$

→ derivatives of (Σ_2) yield coherent results, e.g.,

$$(u_{33})_2 = u_{123} = u_{113} \quad \text{and} \quad (u_{23})_3 = u_{133} = u_{113}$$

→ (Σ_2) is formally integrable with formal power series solutions

$$u_0 + u_i x^i + \frac{u_{11}}{2} (x^1)^2 + u_{11} x^1 x^2 + u_{13} x^1 x^3 + \frac{u_{22}}{2} (x^2)^2 + u_{13} x^2 x^3 + \frac{u_{13}}{2} (x^3)^2 + \dots$$

Integrability defect and homology

Janet example. $(\Sigma) : u_{33} = x^2 u_{11}, \quad u_{22} = 0$

→ $(u_{22})_{33} = 0$ and

$$(u_{33})_{22} = \frac{\partial}{\partial x^2} (x^2 u_{112} + u_{11}) = x^2 u_{1122} + 2u_{112} = 2u_{112}$$

→ new integrability condition $u_{112} = 0$

Remark. The Spencer cohomology does not vanish in homological degree 2

Integrability defect and homology

Janet example. $(\Sigma) : u_{33} = x^2 u_{11}, \quad u_{22} = 0$

→ $(u_{22})_{33} = 0$ and

$$(u_{33})_{22} = \frac{\partial}{\partial x^2} (x^2 u_{112} + u_{11}) = x^2 u_{1122} + 2u_{112} = 2u_{112}$$

→ **new integrability condition** $u_{112} = 0$

Remark. The Spencer cohomology does not vanish in homological degree 2

Integrability defect and homology

Janet example. $(\Sigma) : u_{33} = x^2 u_{11}, \quad u_{22} = 0$

→ $(u_{22})_{33} = 0$ and

$$(u_{33})_{22} = \frac{\partial}{\partial x^2} (x^2 u_{112} + u_{11}) = x^2 u_{1122} + 2u_{112} = 2u_{112}$$

→ new integrability condition $u_{112} = 0$

Remark. The **Spencer cohomology does not vanish in homological degree 2**

Integrability defect and homology

Janet example. $(\Sigma) : u_{33} = x^2 u_{11}, \quad u_{22} = 0$

→ $(u_{22})_{33} = 0$ and

$$(u_{33})_{22} = \frac{\partial}{\partial x^2} (x^2 u_{112} + u_{11}) = x^2 u_{1122} + 2u_{112} = 2u_{112}$$

→ new integrability condition $u_{112} = 0$

Remark. The Spencer cohomology does not vanish in homological degree 2

Theorem (Goldschmidt). If the symbol of Σ is 2-acyclic and if 1st derivatives of Σ do not bring new integrability conditions, then Σ is formally integrable

Integrability defect and homology

Janet example. $(\Sigma) : u_{33} = x^2 u_{11}, \quad u_{22} = 0$

→ $(u_{22})_{33} = 0$ and

$$(u_{33})_{22} = \frac{\partial}{\partial x^2} (x^2 u_{112} + u_{11}) = x^2 u_{1122} + 2u_{112} = 2u_{112}$$

→ new integrability condition $u_{112} = 0$

Remark. The Spencer cohomology does not vanish in homological degree 2

Theorem (Goldschmidt). If the symbol of Σ is 2-acyclic and if 1st derivatives of Σ do not bring new integrability conditions, then Σ is formally integrable

Integrability defect and homology

Janet example. $(\Sigma) : u_{33} = x^2 u_{11}, \quad u_{22} = 0$

→ $(u_{22})_{33} = 0$ and

$$(u_{33})_{22} = \frac{\partial}{\partial x^2} (x^2 u_{112} + u_{11}) = x^2 u_{1122} + 2u_{112} = 2u_{112}$$

→ new integrability condition $u_{112} = 0$

Remark. The Spencer cohomology does not vanish in homological degree 2

Theorem (Goldschmidt). If the symbol of Σ is 2-acyclic and if 1st derivatives of Σ do not bring new integrability conditions, then Σ is formally integrable

Theorem (Serre). Spencer and Koszul complexes are dual to each other

→ 2-acyclicity \Leftrightarrow Koszul homology vanishes in degree 2

Integrability defect and homology

Janet example. $(\Sigma) : u_{33} = x^2 u_{11}, \quad u_{22} = 0$

→ $(u_{22})_{33} = 0$ and

$$(u_{33})_{22} = \frac{\partial}{\partial x^2} (x^2 u_{112} + u_{11}) = x^2 u_{1122} + 2u_{112} = 2u_{112}$$

→ new integrability condition $u_{112} = 0$

Remark. The Spencer cohomology does not vanish in homological degree 2

Theorem (Goldschmidt). If the symbol of Σ is 2-acyclic and if 1st derivatives of Σ do not bring new integrability conditions, then Σ is formally integrable

Theorem (Serre). Spencer and Koszul complexes are dual to each other

→ 2-acyclicity \Leftrightarrow **Koszul homology** vanishes in degree 2

Our contribution:

effectively check 2-acyclicity of linear systems

→ ORE MORPHISMS and ORE MODULES **Maple** packages

Our contribution:

effectively check 2-acyclicity of **linear systems**

→ ORE MORPHISMS and ORE MODULES **Maple** packages

II. COMPUTATION OF KOSZUL HOMOLOGY

Let a system $(\Sigma) : \forall 1 \leq i \leq q, \quad R_1^i u^1 + \cdots + R_p^i u^p = 0$

→ described by $(R_j^i) \in \mathcal{D}^{q \times p}$ (\mathcal{D} : ring of partial differential operators)

Let a system $(\Sigma) : \forall 1 \leq i \leq q, \quad R_1^i u^1 + \cdots + R_p^i u^p = 0$

→ described by $(R_j^i) \in \mathcal{D}^{q \times p}$ (\mathcal{D} : ring of partial differential operators)

Symbol module: $\mathcal{M}(\Sigma) := \mathcal{A}^{1 \times p} / (\mathcal{A}^{1 \times q} \sigma(R))$

→ $\mathcal{A} := \text{gr}(\mathcal{D})$ and $\sigma(R)$: **top-order part** of R

Let a system $(\Sigma) : \forall 1 \leq i \leq q, \quad R_1^i u^1 + \cdots + R_p^i u^p = 0$

→ described by $(R_j^i) \in \mathcal{D}^{q \times p}$ (\mathcal{D} : ring of partial differential operators)

Symbol module: $\mathcal{M}(\Sigma) := \mathcal{A}^{1 \times p} / (\mathcal{A}^{1 \times q} \sigma(R))$

→ $\mathcal{A} := \text{gr}(\mathcal{D})$ and $\sigma(R)$: top-order part of R

Koszul homology: homology $H_k \simeq \ker(\partial_k) / \text{im}(\partial_{k+1})$ of the **Koszul complex**

$$0 \rightarrow \Lambda^n T \otimes \mathcal{M}(\Sigma) \xrightarrow{\partial_n} \cdots \xrightarrow{\partial_3} \Lambda^2 T \otimes \mathcal{M}(\Sigma) \xrightarrow{\partial_2} T \otimes \mathcal{M}(\Sigma) \xrightarrow{\partial_1} \mathcal{M}(\Sigma) \rightarrow 0$$

where for basis elements (χ_i) of $T = \mathcal{A}_1^n$

$$\partial_k (\chi_1 \wedge \cdots \wedge \chi_k \otimes m) := \sum_{i=1}^k (-1)^{i+1} \chi_1 \wedge \cdots \wedge \widehat{\chi}_i \wedge \cdots \wedge \chi_k \otimes \chi_i m$$

Let a system $(\Sigma) : \forall 1 \leq i \leq q, \quad R_1^i u^1 + \cdots + R_p^i u^p = 0$

→ described by $(R_j^i) \in \mathcal{D}^{q \times p}$ (\mathcal{D} : ring of partial differential operators)

Symbol module: $\mathcal{M}(\Sigma) := \mathcal{A}^{1 \times p} / (\mathcal{A}^{1 \times q} \sigma(R))$

→ $\mathcal{A} := \text{gr}(\mathcal{D})$ and $\sigma(R)$: top-order part of R

Koszul homology: homology $H_k \simeq \ker(\partial_k) / \text{im}(\partial_{k+1})$ of the Koszul complex

$$0 \rightarrow \Lambda^n T \otimes \mathcal{M}(\Sigma) \xrightarrow{\partial_n} \cdots \xrightarrow{\partial_3} \Lambda^2 T \otimes \mathcal{M}(\Sigma) \xrightarrow{\partial_2} T \otimes \mathcal{M}(\Sigma) \xrightarrow{\partial_1} \mathcal{M}(\Sigma) \rightarrow 0$$

where for basis elements (χ_i) of $T = \mathcal{A}_1^n$

$$\partial_k(\chi_1 \wedge \cdots \wedge \chi_k \otimes m) := \sum_{i=1}^k (-1)^{i+1} \chi_1 \wedge \cdots \wedge \widehat{\chi}_i \wedge \cdots \wedge \chi_k \otimes \chi_i m$$

Definitions. $\mathcal{M}(\Sigma)$ is **2-acyclic** if H_2 vanishes, and **involutiv** if H_2, \dots, H_n vanish

Objective: compute $H_2 \rightarrow$ using OreMorphisms package

Approach: given $\mathcal{M}_2 \xrightarrow{f} \mathcal{M}_1 \xrightarrow{g} \mathcal{M}_0$ s.t. $g \circ f = 0$ and $\mathcal{M}_i = \mathcal{A}^{1 \times p_i} / (\mathcal{A}^{1 \times q_i} R_i)$

$$\begin{array}{ccccccc}
 \mathcal{A}^{1 \times q_2} & \xrightarrow{\cdot R_2} & \mathcal{A}^{1 \times p_2} & \longrightarrow & \mathcal{M}_2 & \longrightarrow & 0 \\
 \cdot Q_2 \downarrow & & \cdot P_2 \downarrow & & f \downarrow & & \\
 \mathcal{A}^{1 \times q_1} & \xrightarrow{\cdot R_1} & \mathcal{A}^{1 \times p_1} & \longrightarrow & \mathcal{M}_1 & \longrightarrow & 0 \\
 \cdot Q_1 \downarrow & & \cdot P_1 \downarrow & & g \downarrow & & \\
 \mathcal{A}^{1 \times q_0} & \xrightarrow{\cdot R_0} & \mathcal{A}^{1 \times p_0} & \longrightarrow & \mathcal{M}_0 & \longrightarrow & 0
 \end{array}$$

OreMorphisms computes matrices S', S'' s.t. $\ker \begin{pmatrix} \cdot P_1 \\ \cdot R_0 \end{pmatrix} = \mathcal{A}^{1 \times s'} (S' \quad -S'')$ and

$$\ker(g)/\text{im}(f) = \mathcal{A}^{1 \times s'} S' / \left(\mathcal{A}^{1 \times (p_2 + q_2)} \begin{pmatrix} P_2 \\ R_1 \end{pmatrix} \right)$$

Objective: compute $H_2 \rightarrow$ using OreMorphisms package

Approach: given $\mathcal{M}_2 \xrightarrow{f} \mathcal{M}_1 \xrightarrow{g} \mathcal{M}_0$ s.t. $g \circ f = 0$ and $\mathcal{M}_i = \mathcal{A}^{1 \times p_i} / (\mathcal{A}^{1 \times q_i} R_i)$

$$\begin{array}{ccccccc}
 \mathcal{A}^{1 \times q_2} & \xrightarrow{\cdot R_2} & \mathcal{A}^{1 \times p_2} & \longrightarrow & \mathcal{M}_2 & \longrightarrow & 0 \\
 \cdot Q_2 \downarrow & & \cdot P_2 \downarrow & & f \downarrow & & \\
 \mathcal{A}^{1 \times q_1} & \xrightarrow{\cdot R_1} & \mathcal{A}^{1 \times p_1} & \longrightarrow & \mathcal{M}_1 & \longrightarrow & 0 \\
 \cdot Q_1 \downarrow & & \cdot P_1 \downarrow & & g \downarrow & & \\
 \mathcal{A}^{1 \times q_0} & \xrightarrow{\cdot R_0} & \mathcal{A}^{1 \times p_0} & \longrightarrow & \mathcal{M}_0 & \longrightarrow & 0
 \end{array}$$

OreMorphisms computes matrices S', S'' s.t. $\ker \begin{pmatrix} \cdot P_1 \\ \cdot R_0 \end{pmatrix} = \mathcal{A}^{1 \times s'} (S' \quad -S'')$ and

$$\ker(g)/\text{im}(f) = \mathcal{A}^{1 \times s'} S' / \left(\mathcal{A}^{1 \times (p_2 + q_2)} \begin{pmatrix} P_2 \\ R_1 \end{pmatrix} \right)$$

Case of H_2 : $\mathcal{M}_i = \Lambda^i T \otimes \mathcal{M}(\Sigma)$, $f = \partial_{i+1}$, $g = \partial_i$

$\rightarrow R_i = I \otimes \sigma(R)$; $P_i, Q_i = (\text{grad}, \text{curl}, \text{div}) \otimes I$

Theorem (C., Cluzeau, Quadrat). Let (Σ) be a linear system of PDEs and let $\mathcal{M}(\Sigma)$ be the symbol module of (Σ) . Let S, S' be two matrices such that

$$H_2 = \mathcal{A}^{1 \times s'} S' / (\mathcal{A}^{1 \times s} S)$$

Then, $\mathcal{M}(\Sigma)$ is 2-acyclic iff the top degree part of S' is a left-multiple of S .

Theorem (C., Cluzeau, Quadrat). Let (Σ) be a linear system of PDEs and let $\mathcal{M}(\Sigma)$ be the symbol module of (Σ) . Let S, S' be two matrices such that

$$H_2 = \mathcal{A}^{1 \times s'} S' / (\mathcal{A}^{1 \times s} S)$$

Then, $\mathcal{M}(\Sigma)$ is 2-acyclic iff the top degree part of S' is a left-multiple of S .

Algorithmic side of the theorem

Existence of a right factorization
may be checked using OreModules

III. Examples

$(\Sigma) : u_{33} = u_{13}, u_{23} = u_{13}, u_{12} = u_{11}$

Symbol module: $\mathcal{A} := \mathbb{Q}[\chi_1, \chi_2, \chi_3]$

$$\mathcal{M}(\Sigma) = \mathcal{A} / \left(\mathcal{A}^{1 \times 3} \begin{pmatrix} \chi_3^2 - \chi_1 \chi_3 \\ \chi_2 \chi_3 - \chi_1 \chi_3 \\ \chi_1 \chi_2 - \chi_1^2 \end{pmatrix} \right)$$

$$(\Sigma) : u_{33} = u_{13}, u_{23} = u_{13}, u_{12} = u_{11}$$

Symbol module: $\mathcal{A} := \mathbb{Q}[\chi_1, \chi_2, \chi_3]$

$$\mathcal{M}(\Sigma) = \mathcal{A} / \left(\mathcal{A}^{1 \times 3} \begin{pmatrix} \chi_3^2 - \chi_1 \chi_3 \\ \chi_2 \chi_3 - \chi_1 \chi_3 \\ \chi_1 \chi_2 - \chi_1^2 \end{pmatrix} \right)$$

2nd homology group: $H_2 = \mathcal{A}^{1 \times 7} S' / (\mathcal{A}^{1 \times 10} S)$

$$S' = \begin{pmatrix} \chi_3 & \chi_3 & \chi_3 \\ \chi_1 & \chi_2 & \chi_3 \\ 0 & \chi_1 - \chi_2 & 0 \\ 0 & \chi_2 \chi_3 - \chi_3^2 & 0 \\ 0 & 0 & \chi_2 \chi_3 - \chi_3^2 \\ 0 & 0 & \chi_1 \chi_3 - \chi_3^2 \\ 0 & 0 & \chi_1^2 - \chi_1 \chi_2 \end{pmatrix} \quad S = \begin{pmatrix} \chi_1 & \chi_2 & \chi_3 \\ \chi_3^2 - \chi_1 \chi_3 & 0 & 0 \\ \chi_2 \chi_3 - \chi_1 \chi_3 & 0 & 0 \\ \chi_1 \chi_2 - \chi_1^2 & 0 & 0 \\ 0 & \chi_3^2 - \chi_1 \chi_3 & 0 \\ 0 & \chi_2 \chi_3 - \chi_1 \chi_3 & 0 \\ 0 & \chi_1 \chi_2 - \chi_1^2 & 0 \\ 0 & 0 & \chi_3^2 - \chi_1 \chi_3 \\ 0 & 0 & \chi_2 \chi_3 - \chi_1 \chi_3 \\ 0 & 0 & \chi_1 \chi_2 - \chi_1^2 \end{pmatrix}$$

$$(\Sigma) : u_{33} = u_{13}, u_{23} = u_{13}, u_{12} = u_{11}$$

Symbol module: $\mathcal{A} := \mathbb{Q}[\chi_1, \chi_2, \chi_3]$

$$\mathcal{M}(\Sigma) = \mathcal{A} / \left(\mathcal{A}^{1 \times 3} \begin{pmatrix} \chi_3^2 - \chi_1 \chi_3 \\ \chi_2 \chi_3 - \chi_1 \chi_3 \\ \chi_1 \chi_2 - \chi_1^2 \end{pmatrix} \right)$$

2nd homology group: $H_2 = \mathcal{A}^{1 \times 7} S' / (\mathcal{A}^{1 \times 10} S)$

$$S' = \begin{pmatrix} \chi_3 & \chi_3 & \chi_3 \\ \chi_1 & \chi_2 & \chi_3 \\ 0 & \chi_1 - \chi_2 & 0 \\ 0 & \chi_2 \chi_3 - \chi_3^2 & 0 \\ 0 & 0 & \chi_2 \chi_3 - \chi_3^2 \\ 0 & 0 & \chi_1 \chi_3 - \chi_3^2 \\ 0 & 0 & \chi_1^2 - \chi_1 \chi_2 \end{pmatrix} \quad S = \begin{pmatrix} \chi_1 & \chi_2 & \chi_3 \\ \chi_3^2 - \chi_1 \chi_3 & 0 & 0 \\ \chi_2 \chi_3 - \chi_1 \chi_3 & 0 & 0 \\ \chi_1 \chi_2 - \chi_1^2 & 0 & 0 \\ 0 & \chi_3^2 - \chi_1 \chi_3 & 0 \\ 0 & \chi_2 \chi_3 - \chi_1 \chi_3 & 0 \\ 0 & \chi_1 \chi_2 - \chi_1^2 & 0 \\ 0 & 0 & \chi_3^2 - \chi_1 \chi_3 \\ 0 & 0 & \chi_2 \chi_3 - \chi_1 \chi_3 \\ 0 & 0 & \chi_1 \chi_2 - \chi_1^2 \end{pmatrix}$$

2-acyclicity test: **success**

$$(\Sigma) : u_{33} = u_{22} = u_{13} = u_{12} = 0, \quad u_{23} = \alpha u_{11} \quad (\alpha: \text{a parameter})$$

Symbol module: $\mathcal{A} := \mathbb{Q}[\alpha][\chi_1, \chi_2, \chi_3]$

$$\mathcal{M}_\alpha(\Sigma) = \mathcal{A} / \left(\mathcal{A}^{1 \times 5} \begin{pmatrix} \chi_3^2 & \chi_2 \chi_3 - \alpha \chi_1^2 & \chi_2^2 & \chi_1 \chi_3 & \chi_1 \chi_2 \end{pmatrix}^T \right)$$

$$(\Sigma) : u_{33} = u_{22} = u_{13} = u_{12} = 0, \quad u_{23} = \alpha u_{11} \quad (\alpha: \text{a parameter})$$

Symbol module: $\mathcal{A} := \mathbb{Q}[\alpha][\chi_1, \chi_2, \chi_3]$

$$M_\alpha(\Sigma) = \mathcal{A} / \left(\mathcal{A}^{1 \times 5} \begin{pmatrix} \chi_3^2 & \chi_2 \chi_3 - \alpha \chi_1^2 & \chi_2^2 & \chi_1 \chi_3 & \chi_1 \chi_2 \end{pmatrix}^T \right)$$

2nd homology group: $H_2 = \mathcal{A}^{1 \times 11} S' / (\mathcal{A}^{1 \times 16} S)$

$$S' = \begin{pmatrix} \chi_2 & 0 & \alpha \chi_1 \\ \chi_3 & \alpha \chi_1 & 0 \\ \chi_1 & 0 & 0 \\ \chi_3^2 & 0 & 0 \\ \chi_2 \chi_3 & 0 & 0 \\ \chi_2^2 & 0 & 0 \\ 0 & \chi_3 & 0 \\ 0 & \chi_2 & \chi_3 \\ 0 & 0 & \chi_2^2 \\ 0 & 0 & \chi_3^2 \\ 0 & 0 & \chi_1 \chi_3 \end{pmatrix} \quad S = \begin{pmatrix} \chi_1 & \chi_2 & \chi_3 \\ \chi_3^2 & 0 & 0 \\ \chi_2 \chi_3 - \alpha \chi_1^2 & 0 & 0 \\ \chi_2^2 & 0 & 0 \\ \chi_1 \chi_3 & 0 & 0 \\ \chi_1 \chi_2 & 0 & 0 \\ 0 & \chi_3^2 & 0 \\ 0 & \chi_2 \chi_3 - \alpha \chi_1^2 & 0 \\ 0 & \chi_2^2 & 0 \\ 0 & \chi_1 \chi_3 & 0 \\ 0 & \chi_1 \chi_2 & 0 \\ 0 & 0 & \chi_3^2 \\ 0 & 0 & \chi_2 \chi_3 - \alpha \chi_1^2 \\ 0 & 0 & \chi_2^2 \\ 0 & 0 & \chi_1 \chi_3 \\ 0 & 0 & \chi_1 \chi_2 \end{pmatrix}$$

$$(\Sigma) : u_{33} = u_{22} = u_{13} = u_{12} = 0, \quad u_{23} = \alpha u_{11} \quad (\alpha: \text{a parameter})$$

Symbol module: $\mathcal{A} := \mathbb{Q}[\alpha][\chi_1, \chi_2, \chi_3]$

$$M_\alpha(\Sigma) = \mathcal{A} / \left(\mathcal{A}^{1 \times 5} \begin{pmatrix} \chi_3^2 & \chi_2 \chi_3 - \alpha \chi_1^2 & \chi_2^2 & \chi_1 \chi_3 & \chi_1 \chi_2 \end{pmatrix}^T \right)$$

2nd homology group: $H_2 = \mathcal{A}^{1 \times 11} S' / (\mathcal{A}^{1 \times 16} S)$

$$S' = \begin{pmatrix} \chi_2 & 0 & \alpha \chi_1 \\ \chi_3 & \alpha \chi_1 & 0 \\ \chi_1 & 0 & 0 \\ \chi_3^2 & 0 & 0 \\ \chi_2 \chi_3 & 0 & 0 \\ \chi_2^2 & 0 & 0 \\ 0 & \chi_3 & 0 \\ 0 & \chi_2 & \chi_3 \\ 0 & 0 & \chi_2^2 \\ 0 & 0 & \chi_3^2 \\ 0 & 0 & \chi_1 \chi_3 \end{pmatrix} \quad S = \begin{pmatrix} \chi_1 & \chi_2 & \chi_3 \\ \chi_3^2 & 0 & 0 \\ \chi_2 \chi_3 - \alpha \chi_1^2 & 0 & 0 \\ \chi_2^2 & 0 & 0 \\ \chi_1 \chi_3 & 0 & 0 \\ \chi_1 \chi_2 & 0 & 0 \\ 0 & \chi_3^2 & 0 \\ 0 & \chi_2 \chi_3 - \alpha \chi_1^2 & 0 \\ 0 & \chi_2^2 & 0 \\ 0 & \chi_1 \chi_3 & 0 \\ 0 & \chi_1 \chi_2 & 0 \\ 0 & 0 & \chi_3^2 \\ 0 & 0 & \chi_2 \chi_3 - \alpha \chi_1^2 \\ 0 & 0 & \chi_2^2 \\ 0 & 0 & \chi_1 \chi_3 \\ 0 & 0 & \chi_1 \chi_2 \end{pmatrix}$$

2-acyclicity test: **success.**

$$(\Sigma) : u_{33} = u_{22} = u_{13} = u_{12} = 0, \quad u_{23} = \alpha u_{11} \quad (\alpha: \text{a parameter})$$

Symbol module: $\mathcal{A} := \mathbb{Q}[\alpha][x_1, x_2, x_3]$

$$M_\alpha(\Sigma) = \mathcal{A} / \left(\mathcal{A}^{1 \times 5} \begin{pmatrix} x_3^2 & x_2 x_3 - \alpha x_1^2 & x_2^2 & x_1 x_3 & x_1 x_2 \end{pmatrix}^T \right)$$

2nd homology group: $H_2 = \mathcal{A}^{1 \times 11} S' / (\mathcal{A}^{1 \times 16} S)$

$$S' = \begin{pmatrix} x_2 & 0 & \alpha x_1 \\ x_3 & \alpha x_1 & 0 \\ x_1 & 0 & 0 \\ x_3^2 & 0 & 0 \\ x_2 x_3 & 0 & 0 \\ x_2^2 & 0 & 0 \\ 0 & x_3 & 0 \\ 0 & x_2 & x_3 \\ 0 & 0 & x_2^2 \\ 0 & 0 & x_3^2 \\ 0 & 0 & x_1 x_3 \end{pmatrix} \quad S = \begin{pmatrix} x_1 & x_2 & x_3 \\ x_3^2 & 0 & 0 \\ x_2 x_3 - \alpha x_1^2 & 0 & 0 \\ x_2^2 & 0 & 0 \\ x_1 x_3 & 0 & 0 \\ x_1 x_2 & 0 & 0 \\ 0 & x_3^2 & 0 \\ 0 & x_2 x_3 - \alpha x_1^2 & 0 \\ 0 & x_2^2 & 0 \\ 0 & x_1 x_3 & 0 \\ 0 & x_1 x_2 & 0 \\ 0 & 0 & x_3^2 \\ 0 & 0 & x_2 x_3 - \alpha x_1^2 \\ 0 & 0 & x_2^2 \\ 0 & 0 & x_1 x_3 \\ 0 & 0 & x_1 x_2 \end{pmatrix}$$

2-acyclicity test: **success**. Moreover, $M_\alpha(\Sigma)$ is **involution** iff $\alpha \neq 0$

$$(\Sigma) : u_{33} = x^2 u_{11}, \quad u_{22} = 0$$

Symbol module: $\mathcal{A} := \mathbb{Q}[x^1, x^2, x^3][\chi_1, \chi_2, \chi_3]$

$$\mathcal{M}_\alpha(\Sigma) = \mathcal{A} / \left(\mathcal{A}^{1 \times 2} \begin{pmatrix} \chi_3^2 - x^2 \chi_1^2 \\ \chi_2^2 \end{pmatrix} \right)$$

$$(\Sigma) : u_{33} = x^2 u_{11}, \quad u_{22} = 0$$

Symbol module: $\mathcal{A} := \mathbb{Q}[x^1, x^2, x^3][\chi_1, \chi_2, \chi_3]$

$$\mathcal{M}_\alpha(\Sigma) = \mathcal{A} / \left(\mathcal{A}^{1 \times 2} \begin{pmatrix} \chi_3^2 - x^2 \chi_1^2 \\ \chi_2^2 \end{pmatrix} \right)$$

2nd homology group: $H_2 = \mathcal{A}^{1 \times 8} S' / (\mathcal{A}^{1 \times 7} S)$

$$S' = \begin{pmatrix} \chi_1 & \chi_2 & \chi_3 \\ \chi_2^2 & 0 & 0 \\ \chi_2 \chi_3 & 0 & \chi_1 \chi_2 \\ \chi_3^2 & x_2 \chi_1 \chi_2 & x_2 \chi_1 \chi_3 \\ 0 & \chi_2^2 & 0 \\ 0 & x_2 \chi_1^2 - \chi_3^2 & 0 \\ 0 & 0 & \chi_2^2 \\ 0 & 0 & x_2 \chi_1^2 - \chi_3^2 \end{pmatrix} \quad S = \begin{pmatrix} \chi_1 & \chi_2 & \chi_3 \\ \chi_3^2 - x_2 \chi_1^2 & 0 & 0 \\ \chi_2^2 & 0 & 0 \\ 0 & \chi_3^2 - x_2 \chi_1^2 & 0 \\ 0 & \chi_2^2 & 0 \\ 0 & 0 & \chi_3^2 - x_2 \chi_1^2 \\ 0 & 0 & \chi_2^2 \end{pmatrix}$$

$$(\Sigma) : u_{33} = x^2 u_{11}, \quad u_{22} = 0$$

Symbol module: $\mathcal{A} := \mathbb{Q}[x^1, x^2, x^3][\chi_1, \chi_2, \chi_3]$

$$\mathcal{M}_\alpha(\Sigma) = \mathcal{A} / \left(\mathcal{A}^{1 \times 2} \begin{pmatrix} \chi_3^2 - x^2 \chi_1^2 \\ \chi_2^2 \end{pmatrix} \right)$$

2nd homology group: $H_2 = \mathcal{A}^{1 \times 8} S' / (\mathcal{A}^{1 \times 7} S)$

$$S' = \begin{pmatrix} \chi_1 & \chi_2 & \chi_3 \\ \chi_2^2 & 0 & 0 \\ \chi_2 \chi_3 & 0 & \chi_1 \chi_2 \\ \chi_3^2 & x_2 \chi_1 \chi_2 & x_2 \chi_1 \chi_3 \\ 0 & \chi_2^2 & 0 \\ 0 & x_2 \chi_1^2 - \chi_3^2 & 0 \\ 0 & 0 & \chi_2^2 \\ 0 & 0 & x_2 \chi_1^2 - \chi_3^2 \end{pmatrix} \quad S = \begin{pmatrix} \chi_1 & \chi_2 & \chi_3 \\ \chi_3^2 - x_2 \chi_1^2 & 0 & 0 \\ \chi_2^2 & 0 & 0 \\ 0 & \chi_3^2 - x_2 \chi_1^2 & 0 \\ 0 & \chi_2^2 & 0 \\ 0 & 0 & \chi_3^2 - x_2 \chi_1^2 \\ 0 & 0 & \chi_2^2 \end{pmatrix}$$

2-acyclicity test: fail

IV. CONCLUSION AND PERSPECTIVES

Conclusion and perspectives

Summary of presented results

- new criterion for 2-acyclicity of linear PDEs systems
- effective test using `OreMorphisms` and `OreModules` packages
- illustration with various classes of linear systems

Remark. More generally, involutivity can be checked with `OreMorphism` and `OreModules`

Further works

- go further in the effective approach to Spencer cohomology
(Koszul-Tate theory, Spencer sequences)
- applications to physics and control theory
(elasticity theory, hydrodynamics, electromagnetism, general relativity)

Conclusion and perspectives

Summary of presented results

- new criterion for 2-acyclicity of linear PDEs systems
- effective test using `OreMorphisms` and `OreModules` packages
- illustration with various classes of linear systems

Remark. More generally, **involutivity** can be checked with `OreMorphism` and `OreModules`

Further works

- go further in the effective approach to Spencer cohomology
(Koszul-Tate theory, Spencer sequences)
- applications to physics and control theory
(elasticity theory, hydrodynamics, electromagnetism, general relativity)

Conclusion and perspectives

Summary of presented results

- new criterion for 2-acyclicity of linear PDEs systems
- effective test using `OreMorphisms` and `OreModules` packages
- illustration with various classes of linear systems

Remark. More generally, involutivity can be checked with `OreMorphism` and `OreModules`

Further works

- go further in the effective approach to Spencer cohomology
(Koszul-Tate theory, Spencer sequences)
- applications to physics and control theory
(elasticity theory, hydrodynamics, electromagnetism, general relativity)

THANK YOU FOR LISTENING!